


Prostate Prescription Medicine?

What You Don't Know About Prostate Drugs
Can Kill You!



Discover
The
Truth
That Big

Pharmaceutical Companies Don't
Want You To Know

Discover The Truth About The Side Effects Of Prostate Drugs That Major Pharmaceutical Companies Has Been Hiding From You!

Listed below are leading prostate medication that are available in the market today... learn why you must know about the side effects of these drugs.

If you have any question , please feel free to [contact us here](#)

Enjoy your reading.

To your good health

Willie Walter

“How I Solve My Sexual & Prostate Problems Without Viagra Or Any Other Prostate Drugs Within 7 Days! “. If You're Above 40's & Experiencing Low Sex Drive, Frequent Urination, Lack Of Energy & Low Energy Level, [Click Here To READ THIS IMMEDIATELY...](#)

AVODART

“How I Solve My Sexual & Prostate Problems Without Viagra Or Any Other Prostate Drugs Within 7 Days! “. If You're Above 40's & Experiencing Low Sex Drive, Frequent Urination, Lack Of Energy & Low Energy Level, [Click Here To READ THIS IMMEDIATELY...](#)

Avodart

Avodart is a prescription medication used for the treatment of an enlarged prostate. The medicine works by blocking the conversion of testosterone into DHT, which is responsible for enlarging the prostate.

Avodart comes in a capsule form and is taken orally once a day. Among the conditions you should let your healthcare provider know about before you take Avodart are liver disease or failure and any allergies you may have. There is currently no approved generic version of Avodart for sale on the market.

Avodart is manufactured by GlaxoSmithKline.

Avodart works by blocking the conversion of testosterone into DHT (dihydrotestosterone), thereby decreasing the amount of DHT in the body. Since DHT is responsible for enlarging the prostate, using Avodart to decrease DHT levels can treat an enlarged prostate. In most men, Avodart can help shrink the enlarged prostate, helping to relieve their enlarged prostate symptoms.

Avodart Effects

There have been several studies looking at how well Avodart treats an enlarged prostate. These studies showed the following effects related to Avodart and men with BPH.

Symptoms of BPH

In these studies, the men were asked to rate the following symptoms of BPH:

- Incomplete bladder emptying
- Frequent urination
- Intermittent stops and starts in the urine flow
- Feeling of urgency when needing to urinate
- Weak urine stream
- Having to get up to urinate frequently during the night.

The men who took Avodart had significant improvement in their BPH symptoms compared with the men who did not take Avodart. While some patients saw improvement right away, in general it took about 3 to 12 months to see the improvement for most men. This improvement lasted throughout the entire study period (two years).

“How I Solve My Sexual & Prostate Problems Without Viagra Or Any Other Prostate Drugs Within 7 Days! “. If You're Above 40's & Experiencing Low Sex Drive, Frequent Urination, Lack Of Energy & Low Energy Level, [Click Here To READ THIS IMMEDIATELY...](#)

Acute Urinary Retention

These studies also looked at how many men had acute urinary retention. If an [enlarged prostate](#) is very severe, a man may not be able to urinate at all. This is known medically as acute urinary retention and usually requires catheterization (inserting a small tube through the penis into the bladder in order to empty the urine). Of the men who took Avodart, 1.8 percent experienced acute urinary retention that required catheterization, compared to 4.2 percent of men who did not take Avodart.

Surgery

In these studies, 2.2 percent of men who took Avodart needed to have [BPH](#) surgery, while 4.1 percent of men who did not take Avodart needed to have surgery.

Prostate Size

These studies showed that Avodart helped to shrink the prostate gland in most men.

Urine Flow

Men who took Avodart also had an improvement in the rate of urine flow. This increased flow rate was seen as early as the third month after starting Avodart and continued throughout the entire study.

When and How to Take Avodart

Some general considerations for when and how to take Avodart include:

- Avodart comes in a capsule form that is taken by mouth once daily.
- It may take as long as 3 to 12 months of taking Avodart before results can be seen.
- Avodart can be taken with or without food.
- Avodart capsules should be taken whole and should not be chewed or crushed.
- Avodart should be taken at the same time each day to maintain an even level of the drug in your blood.
- For Avodart to work properly, it must be taken as prescribed. Avodart will not work if you stop taking it.

Avodart Dosing

There is only one dose of Avodart for [enlarged prostate treatment](#) -- Avodart 0.5 mg once daily.

As with any medication, do not adjust your Avodart dose unless your healthcare provider specifically instructs you to do so.

“How I Solve My Sexual & Prostate Problems Without Viagra Or Any Other Prostate Drugs Within 7 Days! “. If You're Above 40's & Experiencing Low Sex Drive, Frequent Urination, Lack Of Energy & Low Energy Level, [Click Here To READ THIS IMMEDIATELY...](#)

Side Effects of Avodart

As with any medicine, there are possible side effects with Avodart. However, not everyone who takes Avodart will experience side effects. In fact, most people tolerate Avodart well. When Avodart side effects do occur, they are often minor and either require no treatment or can easily be treated by you or your healthcare provider. Serious side effects are less common.

The most common side effects of Avodart include:

- **Erectile Dysfunction (ED) or Impotence**
- **A decreased libido (decreased sex drive)**
- **Decreased ejaculate amount**
- **Breast enlargement or breast tenderness.**

Caution : Before You Take Avodart

You should talk with your healthcare provider before taking Avodart if you have:

- Liver disease or failure, including cirrhosis
- Any allergies, including allergies to foods, dyes, or preservatives.

Also let your healthcare provider know if you are:

- Pregnant or thinking of becoming pregnant
- Breastfeeding.

Tell your healthcare provider about all of the medicines you currently take, including prescription and non-prescription medicines, vitamins, and herbal supplements.

Overdose With Avodart

In studies, large doses of Avodart did not cause any overdose symptoms.

If you happen to overdose on Avodart, or if you are a pregnant woman who has taken any amount of Avodart, seek medical attention immediately.

“How I Solve My Sexual & Prostate Problems Without Viagra Or Any Other Prostate Drugs Within 7 Days! “. If You're Above 40's & Experiencing Low Sex Drive, Frequent Urination, Lack Of Energy & Low Energy Level, [Click Here To READ THIS IMMEDIATELY...](#)

Avodart Storage Methods

Avodart should be stored at room temperature away from moisture or heat. Avodart capsules should be kept in an airtight container. Because Avodart is so dangerous to pregnant women, women should not take Avodart if they are pregnant or may be pregnant.

Keep Avodart and all medications out of the reach of children

“How I Solve My Sexual & Prostate Problems Without Viagra Or Any Other Prostate Drugs Within 7 Days! “. If You're Above 40's & Experiencing Low Sex Drive, Frequent Urination, Lack Of Energy & Low Energy Level, [Click Here To READ THIS IMMEDIATELY...](#)

Proscar

"How I Solve My Sexual & Prostate Problems Without Viagra Or Any Other Prostate Drugs Within 7 Days! ". If You're Above 40's & Experiencing Low Sex Drive, Frequent Urination, Lack Of Energy & Low Energy Level, [Click Here To READ THIS IMMEDIATELY...](#)

Proscar

Proscar is a prescription drug that can be used for the treatment of benign prostatic hyperplasia, a condition in which the prostate becomes enlarged. By decreasing DHT levels, the medication can help shrink an enlarged prostate and relieve enlarged prostate symptoms.

Proscar comes as a 5 mg tablet and is generally taken once a day. Potential side effects of the medicine may include impotence, decreased sex drive, breast enlargement, and decreased ejaculate amount.

Proscar is manufactured by Merck & Co, Inc. Generic Proscar is made by several manufacturers.

How Does Proscar Work?

Proscar works by blocking the conversion of testosterone into DHT (dihydrotestosterone), which decreases the amount of DHT in the body. Since DHT is important for enlargement of the prostate, using Proscar to decrease DHT levels can treat an enlarged prostate. In most men, Proscar can help shrink the enlarged prostate, helping to relieve their enlarged prostate symptoms

Proscar Effects

There have been several studies looking at how well Proscar treats an enlarged prostate. In particular, there has been one very large study that studied several of the effects of Proscar for men with BPH, with the following results:

Symptoms of BPH

The men in this study were asked to rate the following symptoms of [BPH](#):

- Decreased size and force of urine stream
- Feeling of incomplete bladder emptying when urinating
- Delayed or interrupted urination
- Having to get up to urinate frequently during the night
- Having to urinate frequently during the day
- Needing to strain or push the flow of urine.

The men who took Proscar had significant improvement in their BPH symptoms compared to the men who did not take Proscar. While some patients saw improvement right away, in general, it

"How I Solve My Sexual & Prostate Problems Without Viagra Or Any Other Prostate Drugs Within 7 Days! ". If You're Above 40's & Experiencing Low Sex Drive, Frequent Urination, Lack Of Energy & Low Energy Level, [Click Here To READ THIS IMMEDIATELY...](#)

took about six months to see the improvement for most men. This improvement lasted throughout the entire study period (four years).

Acute Urinary Retention

This study also looked at how many men had acute urinary retention. If an enlarged prostate is very severe, a man may not be able to urinate at all. This is known medically as acute urinary retention and usually requires catheterization (inserting a small tube through the penis into the bladder in order to empty the urine). Of the men who took Proscar, 2.8 percent experienced acute urinary retention that required catheterization, compared to 6.6 percent of men who did not take Proscar.

Surgery

In this study, 4.6 percent of men who took Proscar needed to have BPH surgery, while 10.1 percent of men who did not take Proscar needed to have surgery.

Urine Flow

Men who took Proscar also had an improvement in the rate of urine flow. This increased flow rate was seen as early as the fourth month after starting Proscar and continued throughout the entire study.

When and How to Take Proscar

Some general considerations for when and how to take Proscar include:

- Proscar comes in tablet form. It is taken by mouth once daily.
- It may take as long as six months before results can be seen with Proscar.
- Proscar can be taken with or without food.
- Proscar should be taken at the same time each day to maintain an even level of the drug in your blood.
- For Proscar to work properly, it must be taken as prescribed. Proscar will not work if you stop taking it.

Common Side Effects of Proscar

Proscar has been thoroughly studied in clinical trials, with many people having been evaluated. In these studies, side effects occurring in the group of people taking the drug are always documented and compared to the side effects that occur in a similar group of people not taking the medicine. Based on these studies, the most common Proscar side effects include:

“How I Solve My Sexual & Prostate Problems Without Viagra Or Any Other Prostate Drugs Within 7 Days! “. If You're Above 40's & Experiencing Low Sex Drive, Frequent Urination, Lack Of Energy & Low Energy Level, [Click Here To READ THIS IMMEDIATELY...](#)

- [Erectile dysfunction \(ED\)](#) or [impotence](#) -- in up to 18.5 percent of people
- Decreased libido (sex drive) -- up to 10 percent
- Abnormal ejaculation, including decreased ejaculate amount -- up to 7.2 percent
- Breast enlargement -- up to 2.2 percent.

Proscar Sexual Side Effects: An Overview

Proscar[®] ([finasteride](#)) is a prescription medication that is used for the treatment of an [enlarged prostate](#). As with all medicines, Proscar has possible side effects. Some of these side effects can affect a person's sexual well-being. In the case of Proscar, sexual side effects that have been reported include:

- Decreased sex drive (libido).
- [Impotence](#). Impotence is usually defined as a total inability to achieve erection, an inconsistent ability to do so, or a tendency to sustain only brief erections. Ultimately, impotence is the repeated inability to get or keep an erection firm enough for sexual intercourse. Impotence is also known as [erectile dysfunction](#) (or [ED](#) for short).
- Abnormal ejaculation, including decreased ejaculate amount. In general, decreased ejaculate amount does not affect other aspects of sexual function.

How Common Are Proscar Sexual Side Effects?

Before medicines are approved, they must go through several clinical studies where thousands of people are given a particular medicine and compared to a group of people not given the medicine. In these studies, side effects are always documented. This way, it is possible to see which side effects occur, how often they appear, and how they compare to the group not taking the medicine. Side effects are then usually separated into those that occur in more than 1 percent of people and those that occur in less than 1 percent of people.

With Proscar, the sexual side effects mentioned above occurred in the following percentages of men:

- Erectile dysfunction (ED) or impotence -- in up to 18.5 percent of men
- Decreased libido (sex drive) -- up to 10 percent
- Abnormal ejaculation, including decreased ejaculate amount -- up to 7.2 percent.

Up to 3.7 percent of men stopped taking Proscar because of sexual side effects.

“How I Solve My Sexual & Prostate Problems Without Viagra Or Any Other Prostate Drugs Within 7 Days! “. If You're Above 40's & Experiencing Low Sex Drive, Frequent Urination, Lack Of Energy & Low Energy Level, [Click Here To READ THIS IMMEDIATELY...](#)

Proscar, Sexual Side Effects, and Suggestions

It is impossible for your healthcare provider to know beforehand whether you will develop any sexual side effects while taking [Proscar](#). Also, if a person does develop sexual side effects, such as [impotence](#) or a decrease in libido, it is difficult to know whether these problems are actually caused by another condition or Proscar itself.

However, if you develop any sexual side effects while taking Proscar, make sure to talk with your healthcare provider. There is treatment that he or she may be able to recommend, such as an [erectile dysfunction medication](#). Depending on how severe your sexual side effects are, he or she may also recommend that you adjust your [Proscar dosage](#) or try another medication.

Proscar and Depression

There are several possible side effects of Proscar, and depression does not appear to be one of them. In clinical trials, depression was not reported as a side effect of the medicine. However, this does not mean that the condition cannot occur with the use of Proscar.

Is Depression a Proscar Side Effect?

There are several possible side effects for people taking [Proscar® \(finasteride\)](#). [Depression](#), however, does not appear to be one of them. In clinical studies, depression was not reported as either a common or rare side effect of Proscar.

Keep in mind that just because depression was not seen during clinical studies with Proscar, it does not mean that the medicine cannot cause this side effect. Clinical trials are designed to factor out many possible variables in order to understand whether the medicine works and what its possible side effects are.

Once the medicine is approved, a wider range of people begin taking the medicine and for longer periods of time. As is often the case, this can lead to other side effects being reported. However, once a medicine is approved, there are usually no more studies conducted on it.

This means that certain side effects that occur very rarely may never be reported. Therefore, in some people, it's possible that a medicine may cause a side effect that hasn't been reported before.

So, if you notice any possible depression symptoms while taking Proscar, or if something "just does not seem right," you should talk to your healthcare provider. Your healthcare provider will be able to diagnose and treat the problem.

"How I Solve My Sexual & Prostate Problems Without Viagra Or Any Other Prostate Drugs Within 7 Days! ". If You're Above 40's & Experiencing Low Sex Drive, Frequent Urination, Lack Of Energy & Low Energy Level, [Click Here To READ THIS IMMEDIATELY...](#)

Proscar and Weight Gain

There does not appear to be a link between Proscar and weight gain. In clinical trials, weight gain was not reported as either a common or a rare side effect of Proscar.

Proscar and Weight Gain: An Overview

There are a number of possible side effects with Proscar. Weight gain, however, does not appear to be one of them. This data comes from clinical trials where Proscar was extensively studied and side effects were documented.

Understanding Clinical Trials

Before medicines are approved, they must go through several clinical studies where thousands of people are given a particular medicine and compared to a group of people not given the medicine. In these studies, side effects are always documented. This way, it is possible to see which side effects occur, how often they appear, and how they compare to the group not taking the medicine. Side effects are then usually separated into those that occur in more than 1 percent of people and those that occur in less than 1 percent of people.

For people taking Proscar, weight gain was not reported as either a common or rare side effect.

Proscar and Hair Loss

Many people may be curious about a possible link between Proscar and hair loss treatment. While the active ingredient in Proscar is the same as the active component in Propecia, a medication that treats male pattern baldness, Proscar is not approved for hair loss treatment. Because the dosages for the two medications are different, the drugs are licensed to treat different conditions

Proscar and Hair Loss: An Overview

Proscar is a prescription medication licensed to treat an enlarged prostate(also known medically as benign prostatic hyperplasia, benign prostatic hypertrophy, or **BPH**). **Propecia**[®], which has the same active ingredient as Proscar, is licensed to treat male pattern baldness.

Proscar comes in 5 mg tablets, while Propecia comes in 1 mg tablets. Proscar (5 mg) should be used for BPH symptoms, and Propecia (1 mg) should be used for male pattern baldness.

“How I Solve My Sexual & Prostate Problems Without Viagra Or Any Other Prostate Drugs Within 7 Days! “. If You're Above 40's & Experiencing Low Sex Drive, Frequent Urination, Lack Of Energy & Low Energy Level, [Click Here To READ THIS IMMEDIATELY...](#)

Can I Use Proscar for Hair Loss?

Because Propecia and Proscar have the same active ingredient, many people wonder, "Can I take Proscar for hair loss?" The answer is generally no. Proscar has not been approved for hair loss treatment. In addition, because Proscar contains more finasteride, it has a higher risk of sexual side effects than Propecia. Only Propecia should be used for hair loss.

Proscar and Hair Loss: A Summary

While Proscar does contain the active ingredient that is approved to treat hair loss, it is not the right strength. Taking Proscar instead of Propecia is likely to cause more sexual side effects. While some healthcare providers may suggest cutting Proscar tablets into quarters to use in place of Propecia, this is not an approved use.

Proscar Warnings and Precautions

There are many important Proscar warnings and precautions to be aware of before starting the medication, including serious side effects that may occur. Proscar can potentially cause side effects, including: breast changes; lowered prostate specific antigen levels; or sexual problems such as decreased libido, impotence, or decreased ejaculate amount. You should not take Proscar if you are or may be pregnant or are allergic to any components of the medicine.

Some Proscar Warnings and Precautions

Some precautions and warnings to be aware of include:

- Proscar is considered a pregnancy Category X medication. This means that Proscar is very dangerous for use during pregnancy. Pregnant women (and women who may be pregnant) should not take Proscar and should not come in contact with broken or crushed Proscar tablets. It is not known whether Proscar is passed through breast milk. Therefore, if you are breastfeeding or plan to start breastfeeding, be sure to talk with your healthcare provider about this.
- Before you start Proscar, your healthcare provider should make sure your enlarged prostate symptoms are not caused by another condition, such as Prostate Cancer or bladder problems.
- Men with severely decreased urine flow or men who can only empty very little of their bladder at a time should be closely monitored by their healthcare providers. For these men, Proscar may not be the best option.

"How I Solve My Sexual & Prostate Problems Without Viagra Or Any Other Prostate Drugs Within 7 Days! ". If You're Above 40's & Experiencing Low Sex Drive, Frequent Urination, Lack Of Energy & Low Energy Level, [Click Here To READ THIS IMMEDIATELY...](#)

- Proscar does not prevent prostate cancer. Men can have both BPH and prostate cancer at the same time.
- [Propecia](#) can lower your PSA levels. The PSA (prostate specific antigen) blood test is a common way to screen for prostate cancer. Your healthcare provider will need to adjust your PSA test result (by doubling it) if you are taking Proscar. Your healthcare provider can also use a slightly different blood test (the "percent free PSA" test) that does not need to be adjusted.
- People with liver problems, including liver failure or cirrhosis, should talk to their healthcare provider before taking Proscar. The liver helps to clear Proscar from the body, and Proscar could build up in people with liver problems.
- Proscar can cause sexual side effects, including decreased libido (sex drive), erectile dysfunction ([ED](#) or [impotence](#)), and decreased ejaculate amount. Usually, decreased ejaculate amount does not interfere with sexual function
- Report any breast changes to your healthcare provider right away. Proscar has been reported to cause breast changes (such as breast enlargement, breast tenderness, or tumors).

Who Should Not Take Proscar?

You should not take proscar if you:

- Are allergic to Proscar, [finasteride](#), or any inactive component used to make Proscar. Your healthcare provider or pharmacist has a list of the inactive ingredients used to make Proscar.
- Are or may be pregnant.

Proscar Precautions and Warnings: Summary

There are a number of situations in which Proscar should not be taken, as well as a number of precautions and warnings to be aware of before starting this drug. However, it is possible that not all precautions or warnings for Proscar were discussed in this article. Therefore, you should talk with your healthcare provider about the Proscar warnings and/or precautions that may apply to you.

“How I Solve My Sexual & Prostate Problems Without Viagra Or Any Other Prostate Drugs Within 7 Days! “. If You're Above 40's & Experiencing Low Sex Drive, Frequent Urination, Lack Of Energy & Low Energy Level, [Click Here To READ THIS IMMEDIATELY...](#)

Proscar and Pregnancy

Women who are or may be pregnant should not take Proscar or come in contact with crushed or broken Proscar tablets. This is because the combination of Proscar and pregnancy can put the fetus at risk of complications. The U.S. Food and Drug Administration classifies Proscar as a pregnancy Category X medication.

Is Proscar Safe During Pregnancy?

For women who are pregnant, [Proscar](#)[®] ([finasteride](#)) is very dangerous. This is based on animal studies that looked at the effects of Proscar during pregnancy. Proscar has been given a pregnancy Category X classification based on its risks to an unborn fetus.

Proscar and Pregnancy Category X

The U.S. Food and Drug Administration (FDA) uses a pregnancy category system to classify the possible risks to a fetus when a specific medicine is taken during pregnancy. Pregnancy Category X is given to medicines that show problems to the fetus in animal studies or in humans who have mistakenly taken the medicine. The use of a pregnancy Category X medicine during pregnancy is not recommended.

Proscar should not be used in women who are pregnant or may become pregnant. In fact, women who are or may be pregnant should not touch broken or crushed Proscar tablets.

Proscar, Pregnancy, and Birth Defects

Proscar works by blocking the conversion of testosterone into DHT (dihydrotestosterone), which decreases the amount of DHT in the body. DHT is important for male genital development; if Proscar is taken during pregnancy, it may cause abnormalities to the external genitals of a male fetus.

Can Proscar Be Used by My Partner While I Am Pregnant?

It appears that when Proscar is used by the male partner during pregnancy, no problems with the fetus will occur as a result of Proscar. This is based on data that looked at doses up to 120 times the amount of Proscar found in semen. These studies were conducted in monkeys, a species that closely resembles humans when looking at fetal development. As discussed above, if your partner is using Proscar, it is extremely important that you do not touch any broken or crushed Proscar tablets. Unbroken tablets are safe because they have a protective coating.

Recommendations for Using Proscar During Pregnancy

First of all, [Proscar](#) has not been approved for any use in women. Secondly, taking Proscar during pregnancy can cause birth defects. Therefore, if you are pregnant or thinking about becoming pregnant, let your healthcare provider know immediately. You should not take Proscar if you are or may become pregnant.

“How I Solve My Sexual & Prostate Problems Without Viagra Or Any Other Prostate Drugs Within 7 Days! “. If You're Above 40's & Experiencing Low Sex Drive, Frequent Urination, Lack Of Energy & Low Energy Level, [Click Here To READ THIS IMMEDIATELY...](#)

Caution : Before You Take Proscar

You should talk with your healthcare provider before taking Proscar if you have:

- Liver disease or liver failure, including cirrhosis
- Any allergies, including allergies to foods, dyes, or preservatives.

Tell your healthcare provider about all other medicines you may currently be taking, including prescription and non-prescription medicines, vitamins, and herbal supplements.

Overdose With Proscar

In studies, large doses of Proscar did not cause any overdose symptoms.

If you happen to overdose on Proscar, or if you are a pregnant woman who has taken any amount of Proscar, seek medical attention immediately.

Proscar Storage Methods

Proscar should be stored at room temperature away from moisture or heat. Proscar tablets should be kept in an airtight container. Because Proscar is so dangerous for pregnant women, women should not touch broken or crushed Proscar tablets if they are pregnant or may be pregnant.

Keep Proscar and all medications out of the reach of children.

Proscar Versus Propecia

[Propecia](#)[®], a medication used for treating male pattern baldness, contains the same active ingredient as Proscar -- [finasteride](#). However, Propecia is available only in 1 mg tablets, which is not the correct dose for treating BPH

Generic Proscar

Generic Proscar is currently licensed for sale.

“How I Solve My Sexual & Prostate Problems Without Viagra Or Any Other Prostate Drugs Within 7 Days! “. If You're Above 40's & Experiencing Low Sex Drive, Frequent Urination, Lack Of Energy & Low Energy Level, [Click Here To READ THIS IMMEDIATELY...](#)

FLOMAX

“How I Solve My Sexual & Prostate Problems Without Viagra Or Any Other Prostate Drugs Within 7 Days! “. If You're Above 40's & Experiencing Low Sex Drive, Frequent Urination, Lack Of Energy & Low Energy Level, [Click Here To READ THIS IMMEDIATELY...](#)

What Is Flomax?

Flomax® (tamsulosin hydrochloride) is a prescription medicine that has been licensed as treatment for an enlarged prostate. An enlarged prostate is also known medically as benign prostatic hyperplasia, benign prostatic hypertrophy, or BPH

Flomax is manufactured by Boehringer Ingelheim Pharmaceuticals, Inc.

Flomax is part of a class of drugs known as alpha blockers. Flomax works by blocking alpha-1 receptors. Alpha-1 receptors are found in several places of the body, including the prostate gland.

For the treatment of BPH, blocking alpha-1 receptors helps to relax the muscles of the prostate and the bladder neck (where urine leaves the bladder). Flomax does not shrink the prostate, as some other BPH medications do. However, by relaxing the muscles of the prostate and bladder, Flomax helps to quickly relieve BPH symptoms. While Flomax can help with symptoms, it is not a cure for BPH.

Flomax Effects

There have been several studies looking at how well Flomax treats enlarged prostate symptoms

Symptoms of BPH

In studies of Flomax for BPH treatment, men were asked to rate the following symptoms of BPH:

- Incomplete bladder emptying
- Frequent urination
- Intermittent stops and starts in the urine flow
- Feeling of urgency when needing to urinate
- Weak urine stream
- Having to get up to urinate frequently during the night.

The men who took Flomax had significant improvement in their BPH symptoms compared with the men who did not take Flomax. This improvement was usually seen within the first week and lasted throughout the study period (13 weeks).

“How I Solve My Sexual & Prostate Problems Without Viagra Or Any Other Prostate Drugs Within 7 Days! “. If You're Above 40's & Experiencing Low Sex Drive, Frequent Urination, Lack Of Energy & Low Energy Level, [Click Here To READ THIS IMMEDIATELY...](#)

Urine Flow

Men who took Flomax also had an improvement in the rate of urine flow. This increased flow rate was seen as early as the first week after starting BPH treatment with Flomax and continued throughout the entire study.

When and How to Take Flomax

Some general considerations for when and how to take Flomax include:

- Flomax comes in capsule form. Flomax is taken by mouth once daily, thirty minutes after a meal.
- Flomax capsules should be taken whole and should not be opened, chewed, or crushed.
- Flomax should be taken at the same time each day to maintain an even level of the drug in your blood.

For Flomax to work properly, it must be taken as prescribed. Flomax will not work if you stop taking it.

Flomax Dosing

When first starting Flomax (or if you have been off Flomax for several days), the dose of Flomax is 0.4 mg once daily. If Flomax does not seem to be working after two to four weeks at this dose, your healthcare provider may increase your dose to 0.8 mg once daily.

As with any medication, do not adjust your Flomax dose unless your healthcare provider specifically instructs you to do so.

Side Effects of Flomax

As with any medicine, there are possible side effects with Flomax. However, not everyone who takes Flomax will experience side effects. In fact, most people tolerate Flomax well. When Flomax side effects do occur, they are often minor and either require no treatment or can easily be treated by you or your healthcare provider. Serious side effects are less common.

The most common side effects of Flomax include:

- Abnormal ejaculation
- Runny or stuffy nose
- Dizziness
- Infections, including the cold or flu.

“How I Solve My Sexual & Prostate Problems Without Viagra Or Any Other Prostate Drugs Within 7 Days! “. If You're Above 40's & Experiencing Low Sex Drive, Frequent Urination, Lack Of Energy & Low Energy Level, [Click Here To READ THIS IMMEDIATELY...](#)

Drug Interactions With Flomax

Flomax can potentially interact with a number of medicines.

Caution : Before You Take Flomax

You should talk with your healthcare provider before taking Flomax if you have:

- Any allergies, including allergies to foods, dyes, or preservatives, especially a sulfa allergy.

Also, let your healthcare provider know if you are:

- Pregnant or thinking of becoming pregnant Breastfeeding.

Tell your healthcare provider about all of the medicines you currently take, including prescription and non-prescription medicines, vitamins, and herbal supplements.

Overdose With Flomax

People who take too much Flomax may have overdose symptoms that can include:

- Low blood pressure (hypotension)
- Headache.

If you happen to overdose on Flomax, seek medical attention immediately.

Keep Flomax and all medications out of the reach of children.

Flomax: Strengths

Flomax capsules are available in one strength -- Flomax 0.4 mg capsules.

Generic Flomax

There is currently no generic Flomax available for sale.

“How I Solve My Sexual & Prostate Problems Without Viagra Or Any Other Prostate Drugs Within 7 Days! “. If You're Above 40's & Experiencing Low Sex Drive, Frequent Urination, Lack Of Energy & Low Energy Level, [Click Here To READ THIS IMMEDIATELY...](#)

CARDURA

“How I Solve My Sexual & Prostate Problems Without Viagra Or Any Other Prostate Drugs Within 7 Days! “. If You're Above 40's & Experiencing Low Sex Drive, Frequent Urination, Lack Of Energy & Low Energy Level, [Click Here To READ THIS IMMEDIATELY...](#)

Cardura

Cardura is commonly used to treat symptoms of an enlarged prostate and to lower high blood pressure. It works by relaxing the muscles of the prostate and bladder. At the same time, it relaxes the blood vessels in the body to help lower blood pressure. Cardura is available in tablet form and is typically taken once a day. Potential side effects include dizziness, drowsiness, and fatigue.

Cardura® (doxazosin mesylate) is a prescription medicine that has been licensed for the following uses:

- Treating an enlarged prostate
- Controlling high blood pressure

Cardura is made by Pfizer, Inc. Several manufacturers make generic Cardura

Cardura is part of a class of drugs known as alpha blockers. The drug works by blocking alpha-1 receptors. These receptors are found in several places of the body and have many different effects.

When used to treat benign prostatic hyperplasia, blocking alpha-1 receptors helps to relax the muscles of the prostate and the bladder neck (where urine leaves the bladder). Cardura does not shrink the prostate, however, as some other BPH medications do. By relaxing the muscles of the prostate and bladder, Cardura helps to quickly relieve BPH symptoms.

When used to treat high blood pressure, blocking alpha-1 receptors helps to relax the blood vessels throughout the body. Tightly constricted blood vessels can be one of the causes of high blood pressure. By relaxing the vessels, they become wider and allow the blood to flow more easily. This decreases the pressure in the vessels.

Effects of Cardura

Several studies have looked at how well Cardura treats enlarged prostate symptoms and high blood pressure.

Benign Prostatic Hyperplasia

In studies on Cardura and benign prostatic hyperplasia treatment, men experienced relief from their symptoms within one week of starting Cardura. Also, the men who took Cardura in these studies had an improvement in their rate of urine flow.

“How I Solve My Sexual & Prostate Problems Without Viagra Or Any Other Prostate Drugs Within 7 Days! “. If You're Above 40's & Experiencing Low Sex Drive, Frequent Urination, Lack Of Energy & Low Energy Level, [Click Here To READ THIS IMMEDIATELY...](#)

High Blood Pressure

When Cardura was studied as a treatment for high blood pressure, people who took the drug lowered their blood pressure more than people who did not take it. This lower blood pressure was seen as soon as one day after taking Cardura.

A blood pressure reading consists of two numbers -- for example, 120/80. The top number is known as the systolic blood pressure, and the bottom number is the diastolic blood pressure. During clinical studies of people taking Cardura, systolic blood pressure decreased 9 to 10 mmHg (millimeters of mercury) on average and diastolic blood pressure decreased by 5 to 8 mmHg on average. By lowering blood pressure, Cardura can decrease the risks that often accompany long-term high blood pressure. However, Cardura is not a cure for high blood pressure.

When and How to Take Cardura

General considerations for when and how to take Cardura include the following:

- Cardura comes in tablet form and is taken by mouth once a day.
- Cardura should be taken at the same time each day to maintain an even level of the drug in your blood.
- You can take Cardura with or without food.

For Cardura to work properly, it must be taken as prescribed. Cardura will not work if you stop taking it.

Dosing Information for Cardura

The dose of Cardura that your healthcare provider recommends will vary, depending on a number of factors, including:

- Your age
- The medical condition you are being treated for
- Other medical conditions you may have
- Other medications you are currently taking
- How you respond to Cardura.

“How I Solve My Sexual & Prostate Problems Without Viagra Or Any Other Prostate Drugs Within 7 Days! “. If You're Above 40's & Experiencing Low Sex Drive, Frequent Urination, Lack Of Energy & Low Energy Level, [Click Here To READ THIS IMMEDIATELY...](#)

Side Effects of Cardura

As with any medicine, side effects are possible with Cardura. However, not everyone who takes the drug will experience side effects. In fact, most people tolerate Cardura quite well. If side effects do occur, they are often minor and either require no treatment or can easily be treated by you or your healthcare provider. Serious side effects are less common.

The most common side effects of Cardura include:

- Dizziness
- Tiredness (fatigue) or a general ill feeling (malaise)
- Sleepiness or drowsiness.

Drug Interactions With Cardura

Cardura can potentially interact with a number of medicines

Caution : Before You Take Cardura

You should talk with your healthcare provider before taking Cardura if you have:

- Liver problems, including liver failure or cirrhosis
- Any allergies, including allergies to foods, dyes, or preservatives.

Also let your healthcare provider know if you are:

- Pregnant or thinking of becoming pregnant
- Breastfeeding.

Tell your healthcare provider about all of the medicines you currently take, including prescription and non-prescription medicines, vitamins, and herbal supplements.

What If I Take an Overdose of Cardura?

People who take too much Cardura may have overdose symptoms that can include:

- Low [blood pressure](#) (hypotension)
- Lightheadedness, dizziness, or fainting
- Drowsiness
- Loss of life.

“How I Solve My Sexual & Prostate Problems Without Viagra Or Any Other Prostate Drugs Within 7 Days! “. If You're Above 40's & Experiencing Low Sex Drive, Frequent Urination, Lack Of Energy & Low Energy Level, [Click Here To READ THIS IMMEDIATELY...](#)

If you happen to overdose on Cardura, seek immediate medical attention.

Keep Cardura and all medications out of the reach of children.

What Should I Do If I Miss a Dose of Cardura?

If you do not take your Cardura as scheduled, take your next dose as soon as you remember. If it is almost time for your next dose, skip the missed dose and continue with your regular schedule. Do not take a double dose of Cardura.

If you miss several doses of Cardura in a row, contact your healthcare provider before restarting Cardura, as you will likely need to restart at a lower dose to prevent side effects.

Strengths of Cardura

Cardura tablets are available in four strengths:

- Cardura 1 mg tablets
- Cardura 2 mg tablets
- Cardura 4 mg tablets
- Cardura 8 mg tablets.

Generic Cardura

There is currently a generic version of Cardura licensed for sale (see [Generic Cardura](#) for more information).

Articles taken from : <http://prostate.emedtv.com/>

“How I Solve My Sexual & Prostate Problems Without Viagra Or Any Other Prostate Drugs Within 7 Days! “. If You're Above 40's & Experiencing Low Sex Drive, Frequent Urination, Lack Of Energy & Low Energy Level, [Click Here To READ THIS IMMEDIATELY...](#)